

We study the gamma noise measure μ on Borel sets of Schwartz space $S'(\mathbb{R})$, which turns out to be an interesting example of compound Poisson measure. The orthogonal analysis with respect to μ relates to biorthogonal one analogously to the case of Poisson measure. This fact enables to solve stochastic Wick–Skorokhod equations involving gamma noise.

1. Let $N' \supset \mathcal{H} \supset N$ be a nuclear rigging of the real Hilbert space $\mathcal{H} = L_{2,Re}(\mathbb{R})$ by the real Schwartz space $N := S(\mathbb{R}) = \text{pr} \lim_{p \in \mathbb{N}} \mathcal{H}_p$ and $N' := S'(\mathbb{R}) = \text{ind} \lim_{p \in \mathbb{N}} \mathcal{H}_{-p}$. Here the family $\{\mathcal{H}_p, p \in \mathbb{N}\}$ of the Hilbert spaces satisfies the condition: $\forall p \in \mathbb{N} \exists p' \in \mathbb{N} : \mathcal{H}_{p'} \hookrightarrow \mathcal{H}_p$ and the embedding belongs to Hilbert–Schmidt class; $\mathcal{H}_{-p}$ is the negative space of the chain $\mathcal{H}_{-p} \supset \mathcal{H} \supset \mathcal{H}_p$ (see, e.g., [2]). We denote by $|\cdot|$ and $|\cdot|_p$ the norms in $\mathcal{H}$ and $\mathcal{H}_p, p \in \mathbb{Z} \setminus \{0\}$, respectively and by $\langle \cdot, \cdot \rangle$ the dual pairings between $\mathcal{H}_p$ and $\mathcal{H}_{-p}$ and between N and N' , given by the extension of the inner product in $\mathcal{H}$ and preserve these notations for complexifications and tensor powers of abovementioned spaces.

We define the gamma noise measure μ on $(N', \mathcal{B}(N'))$ by its characteristic functional

$$C_\mu(\theta) = \int_{N'} \exp\{i\langle x, \theta \rangle\} \mu(dx) = \exp\{-\langle 1, \log(1 - i\theta) \rangle\}, \quad \theta \in N \quad (1)$$

via the Bochner–Minlos theorem. This name is natural for μ , because C_μ coincides with the characteristic functional of the gamma noise, i.e. generalized stochastic process with independent values $\xi' = \{\xi'_t, t \in \mathbb{R}\}$ being the distributional derivative of gamma process ξ (for more details on generalized stochastic processes see [5], on gamma process—[13]).

Indeed, setting in (1) $\theta = \lambda \chi_{[0 \wedge t, 0 \vee t]}$, $\lambda \in \mathbb{R}$, $t \in \mathbb{R} \setminus \{0\}$ one obtain $C_\mu(\theta) = (1 - i\lambda)^{-|t|}$. This function coincides with Fourier transform of the density $p_{|t|}(s) = s^{|t|-1} e^{-s} \chi_{(0, \infty)}(s) / \Gamma(|t|)$, $s \in \mathbb{R}$ of $\Gamma(|t|)$ -distribution. Consider the probability space $(N', \mathcal{B}(N'), \mu)$. The random variable $\xi_t(x) := \langle x, \chi_{[0 \wedge t, 0 \vee t]} \rangle$ has $\Gamma(|t|)$ distribution, so the family of random variables $\xi = \{\xi_t, t \in \mathbb{R}\}$ ($\xi_0 := 0$) is gamma process.

The gamma process is a separable stochastic process with independent increments, so the characteristic function $\mathbf{E} \exp\{i\lambda(\xi_{t_2} - \xi_{t_1})\}$, $t_1 < t_2$ of its increment has Lévy–Khinchine representation (see, e.g., [6,10]). By (1) this representation has the following form

$$\int_{N'} \exp\{i\lambda \langle x, \chi_{[t_1, t_2]} \rangle\} \mu(dx) = \exp\left\{(t_2 - t_1) \int_0^\infty (e^{iu\lambda} - 1) \frac{e^{-u}}{u} du\right\}. \quad (2)$$

In other words, the equality (2) means that the Lévy triple (a, σ^2, β) of the gamma process ξ coincides with

$$a = \int_0^\infty \frac{e^{-u}}{1+u^2} du, \quad \sigma^2 = 0, \quad \beta(\Delta) = \int_{\Delta \cap (0, \infty)} \frac{e^{-u}}{u} du.$$

Thus, the gamma process is a non-decreasing compound Poisson process such that almost every sample path has infinitely many jump discontinuities in each open interval and is locally bounded (see, e.g., [6]). This fact, equality (1) and some results from [13, ch. 3] enables us to describe some sets of full μ -measure. Below we denote by $\mathfrak{D}(\mathbb{R})$ the collection of all the locally countable sets $\mathfrak{d} \subset \mathbb{R}$.

Theorem 1. 1) There exists $p \in \mathbb{R}$ such that $\mu(\mathcal{H}_{-p}) = 1$.
 2) The set

$$M := \left\{ \sum_{t_k \in \mathfrak{D}} h_k \delta_{t_k} : h_k > 0, \mathfrak{d} \in \mathfrak{D}(\mathbb{R}) \right\} \cap C_0(\mathbb{R})' \subset N'$$

has the full μ -measure.

3) Denote for $y \in M$ $y(\tau) = \langle y, \chi_{[0 \wedge \tau, 0 \vee \tau]} \rangle$. Then

- (i) $\forall \sigma > 0 \mu(\{y \in M : |y(\tau) - \tau| \geq \sigma\}) \rightarrow 0, \tau \rightarrow \pm\infty$;
- (ii) $\mu(\{y \in M : y(\tau) - \tau, \tau - y(\tau) \text{ are bounded on } \mathbb{R}\}) = 0$.

Remark. The gamma noise measure is a concrete example of generalized white noise measure. The latter measures were introduced in [5], see also [1,4]. The first statement of Theorem 1 is valid for all the generalized white noise measures (it is a simple consequence of Lévy–Khinchine representation). The second one holds for compound Poisson measures such that the measure β in Lévy triple satisfies the following conditions: β is an infinite measure on $(0, \infty)$; $\forall \delta > 0 \beta((\delta, \infty)) < \infty$; $m_1(\beta) = \int_0^\infty u d\beta(u) < \infty$. The third statement holds for compound Poisson measures, satisfying additional requirement: $m_1(\beta) = 1$.

2. It follows from (1) that the gamma noise measure μ is analytic one on $(N', \mathcal{B}(N'))$. Then the function

$$e_\mu(x; \theta) = \exp\{\langle x, \theta \rangle + \langle 1, \log(1 - \theta) \rangle\}, \quad x \in N', \quad \theta \in N$$

generates the system $\mathbf{P}^\mu$ of Appell polynomials, corresponding to μ , i.e. $e_\mu(x; \theta) = \sum_{n=0}^\infty \frac{\langle P_n^\mu(x), \theta^{\otimes n} \rangle}{n!}$ (see [12]). Let a vector-function $N_{\mathbb{C}} \ni \theta \mapsto \alpha(\theta) \in N_{\mathbb{C}}$ be holomorphic at zero. Then

$$e_\mu^\alpha(x; \theta) := e_\mu(x; \alpha(\theta)), \quad x \in N', \quad \theta \in N$$

is generating function of the system $\mathbf{P}^{\mu, \alpha}$ of generalized Appell polynomials, corresponding to measure μ and function α (for more details see [11], in one-dimensional case $N' = \mathcal{H} = N = \mathbb{R}$ see [3]).

Let π be Poisson measure on $(N', \mathcal{B}(N'))$. Recall that the generalized Appell polynomials $\mathbf{P}^{\pi, \alpha_\pi}$ corresponding to π and $\alpha_\pi(\theta) = \log(1 + \theta)$ coincide with Charlier polynomials, that are orthogonal with respect to π , see [14,15]. In the case of gamma noise measure the following statement is true (see [7,8]).

Theorem 2. Let $\alpha_\mu(\theta) = \frac{\theta}{\theta-1}$, $\theta \in N$. The generalized Appell polynomials $\mathbf{P}^{\mu, \alpha_\mu}$ are orthogonal with respect to gamma noise measure μ .

In one-dimensional case $\mathbf{P}^{\mu, \alpha_\mu}$ coincides with the system of Laguerre polynomials (see, e.g., [3]). So we can call the polynomials from the system $\mathbf{J}^\mu := \mathbf{P}^{\mu, \alpha_\mu}$ infinite-dimensional Laguerre polynomials. As in Gaussian and Poissonian analysis one can construct the rigging $(N)_{\mathbf{J}^\mu}^{-1} \supseteq L_2(N', \mu) \supseteq (N)_{\mathbf{J}^\mu}^1$, using the system $\mathbf{J}^\mu = \{\langle J_n^\mu(\cdot), \varphi^{(n)} \rangle, \varphi^{(n)} \in N_{\mathbb{C}}^{\widehat{\otimes} n}, n \in \mathbb{Z}_+\}$ ($\widehat{\otimes}$ denotes symmetric tensor product) of Laguerre polynomials. Namely, $(N)_{\mathbf{J}^\mu}^1 = \text{pr} \lim_{p, q \in \mathbb{N}} (\mathcal{H}_p)_q^1$, $(N)_{\mathbf{J}^\mu}^{-1} = \text{ind} \lim_{p, q \in \mathbb{N}} (\mathcal{H}_{-p})_q^{-1}$, where $(\mathcal{H}_p)_q^1$ is a completion of the space of continuous polynomials on N' $\mathcal{P}(N') = \{\varphi(x) = \sum_{k=0}^n \langle J_k^\mu(x), \varphi^{(k)} \rangle, \varphi^{(k)} \in N_{\mathbb{C}}^{\widehat{\otimes} k}, k = \overline{0, n}, n \in \mathbb{Z}_+\}$

$\mathbb{Z}_+\}$ with respect to the Hilbert norm $\|\varphi\|_{p,q,1}^2 = \sum_{n=0}^{\infty} (n!)^2 2^{qn} |\varphi^{(n)}|_p^2$, $\varphi \in \mathcal{P}(N')$, $(\mathcal{H}_{-p})_{-q}^{-1}$ is dual of $(\mathcal{H}_p)_q^1$ with respect to $L_2(N', \mu)$.

On another hand, using $\mathbf{J}^\mu$ one can prove a nondegeneracy of the gamma noise measure μ and construct the dual Appell system $\mathbf{Q}^\mu$ of distributions on N' and the rigging $(N)_\mu^{-1} \supset L_2(N', \mu) \supset (N)^1$ of $L_2(N', \mu)$ by Kondratiev spaces of distributions and test functions respectively (see [12]). As in the case of Poisson measure π the equalities $(N)_\mu^{-1} = (N)_{\mathbf{J}^\mu}^{-1}$, $(N)^1 = (N)_{\mathbf{J}^\mu}^1$ are valid, so we have a description of $(N)_{\mathbf{J}^\mu}^{-1}$ and $(N)_{\mathbf{J}^\mu}^1$ via characterization theorems from [12].

We apply this result for solving of the gamma counterpart of Verhulst type equation

$$Y_t = Y_0 + r \int_0^t Y_s \diamond (1 - Y_s) ds + a \int_0^t Y_s \diamond (1 - Y_s) \diamond \xi'_s ds, \quad (3)$$

modelling the population growth in a crowded stochastic environment. Here $\{Y_t, t \in \mathbb{R}_+\} \subset (N)_\mu^{-1}$ is unknown generalized stochastic process, $r \in \mathbb{R}$, $a > 0$, $\diamond$ means Wick product (for more details on this and more general Wick–Skorokhod equations see [9], Poisson counterpart of (3) was considered in [15]).

Theorem 3. *Let $Y_0 \in (N)_\mu^{-1}$ be such that $\langle\langle Y_0, 1 \rangle\rangle \neq 0$ ($\langle\langle \cdot, \cdot \rangle\rangle$ denotes the dual pairing between $(N)_\mu^{-1}$ and $(N)^1$, given by inner product in $L_2(N', \mu)$). Then*

$$Y_t(x) = [1 + (Y_0^{\diamond(-1)} - 1) \diamond e_\mu^{\alpha\mu}(-r - ax; \chi_{[0,t]})]^{\diamond(-1)}, \quad x \in N'$$

is the unique $(N)_\mu^{-1}$ -solution to (3).

The proof uses $S_\mu^{\alpha\mu}$ -transform of $\Phi \in (N)_\mu^{-1}$: $(S_\mu^{\alpha\mu} \Phi)(\theta) = \langle\langle \Phi, e_\mu^{\alpha\mu}(\cdot; \theta) \rangle\rangle$, being the analogue of S -transform of Gaussian and S_π -transform of Poissonian analysis. Namely, we have for any $\Phi, \Psi \in (N)_\mu^{-1} = (N)_{\mathbf{J}^\mu}^{-1}$ that $S_\mu^{\alpha\mu}(\Phi \diamond \Psi)(\theta) = S_\mu^{\alpha\mu}(\Phi)(\theta) S_\mu^{\alpha\mu}(\Psi)(\theta)$, $\theta \in N$. So $S_\mu^{\alpha\mu}$ transfers the equation (3) to deterministic one as well as S and S_π . It remains to apply the standard arguments of white noise functional approach to Wick–Skorokhod equations [9].

REFERENCES

1. S. Albeverio, J.-L. Wu, *Euclidean Random Fields Obtained by Convolution from Generalized White Noise*, SFB237—Preprint No. 245, Ruhr–Universität–Bochum (1995).
2. Yu. M. Berezansky, Yu. G. Kondratiev, *Spectral Methods in Infinite Dimensional Analysis*, Naukova Dumka, Kiev (1988); English transl. Kluwer AP, Dordrecht (1994).
3. R. P. Boas, R. C. Buck, *Polynomial Expansions of Analytic Functions*, Springer, Berlin, 1964.
4. A. Dermoune, Ann. IHP, Prob. et Stat. **26**, **1** (1990), pp. 101–119.
5. I. M. Gelfand, N. Ya. Vilenkin, *Generalized Function IV. Some Applications of Harmonic Analysis*, Fizmatgiz, Moscow (1961); English transl. AP, New York (1964).
6. I. I. Gihman, A. V. Skorokhod, *Theory of Stochastic Processes*. I, Nauka, Moscow, 1971. (in Russian)
7. A. V. Gorbunov, *Dual Appell System Corresponding to Gamma Noise*, In: Proceedings of Ukrainian Conference of Young Scientists (Mathematics), Kiev University (1994), pp. 87–97. (in Russian)
8. A. V. Gorbunov, *Non-Gaussian Infinite-Dimensional Analysis in the Case of Gamma-Noise (Biorthogonal Approach)*, In: Conference on Stochastic Differential and Differential Equations, Győr, Hungary, August 21–24, 1996, Abstracts (1996), pp. 103–104.
9. H. Holden, B. Oksendal, J. Ubøe and T. S. Zhang, *Stochastic Partial Differential Equations*, Oslo, 1996.

10. K. Itô, Proc AMS **81**, **2** (1956), pp. 253–263.
11. Yu. G. Kondratiev, J. Luis da Silva, L. Streit, *Generalized Appell Systems*, BiBoS preprint 729/5/96 (1996).
12. Yu. G. Kondratiev, L. Streit, W. Westerkamp, J. Yan, *Generalized Functions in Infinite Dimensional Analysis*, IAS Reports No. 1995–002, Kyoto (1995).
13. L. Takacz, *Combinatorial Methods in the Theory of Stochastic Processes*, John Wiley & Sons, New York, 1967.
14. G. F. Us, Methods Funct. Anal. Topol. **v.1**, **No. 1** (1995), pp. 93–108.
15. G. F. Us, *One Application of Poisson Infinite Dimensional Analysis*, Proc. of Crimea math. school - VI (1996), 319.

A. V. GORBUNOV, ASSISTANT, DEPARTMENT OF ECONOMICS, KIEV UNIVERSITY, VOLODYMYRSKA ST., 64, KIEV, 252033, UKRAINE.

G. F. US, PH.D., ASSOCIATE PROFESSOR, DEPARTMENT OF MECHANICS AND MATHEMATICS, KIEV UNIVERSITY, VOLODYMYRSKA ST., 64, KIEV, 252033, UKRAINE.